

BeppoSAX AND *Chandra* OBSERVATIONS OF SAX J0103.2–7209 = 2E 0101.5–7225: A NEW PERSISTENT 345 S X–RAY PULSAR IN THE SMC[†]G.L. ISRAEL^{1,*}, S. CAMPANA^{2,*}, S. COVINO², D. DAL FIUME³, T.J. GAETZ⁴, S. MEREGHETTI⁵, T. OOSTERBROEK⁶, M. ORLANDINI³, A.N. PARMAR⁶, D. RICCI⁷ AND L. STELLA^{1,*}

1. Osservatorio Astronomico di Roma, Via Frascati 33, I–00040 Monteporzio Catone (Roma), Italy

2. Osservatorio Astronomico di Brera, Via Bianchi 46, I–23807 Merate (Lc), Italy

3. TeSRE, C.N.R., Via Gobetti 101, I–40129 Bologna, Italy

4. Harvard–Smithsonian Center for Astrophysics, MS–70, 60 Garden Street, Cambridge, MA 02138, USA

5. IFCTR, C.N.R., Via Bassini 15, I–20133 Milano, Italy

6. Astrophysics Division, Space Science Department of ESA, ESTEC, P.O. Box 299, 2200 AG Noordwijk, The Netherlands

7. SAX Science Data Center, ASI, Viale Regina Margherita 202, I–00198 Roma, Italy

*Received: 1999 December 6; Accepted: 2000 January 24***ABSTRACT**

We report the results of a 1998 July *BeppoSAX* observation of a field in the Small Magellanic Cloud (SMC) which led to the discovery of ~ 345 s pulsations in the X–ray flux of SAX J0103.2–7209. The *BeppoSAX* X–ray spectrum is well fit by an absorbed power–law with photon index ~ 1.0 plus a black body component with $kT = 0.11$ keV. The unabsorbed luminosity in the 2–10 keV energy range is $\sim 1.2 \times 10^{36}$ erg s^{–1}. In a very recent *Chandra* observation the 345 s pulsations are also detected. The available period measurements provide a constant period derivative of -1.7 s yr^{–1} over the last three years making SAX J0103.2–7209 one of the most rapidly spinning–up X–ray pulsars known. The *BeppoSAX* position (30'' uncertainty radius) is consistent with that of the *Einstein* source 2E 0101.5–7225 and the *ROSAT* source RX J0103.2–7209. This source was detected at a luminosity level of few 10^{35} – 10^{36} erg s^{–1} in all datasets of past X–ray missions since 1979. The *ROSAT* HRI and *Chandra* positions are consistent with that of a $m_V = 14.8$ Be spectral type star already proposed as the likely optical counterpart of 2E 0101.5–7225. We briefly report and discuss photometric and spectroscopic data carried out at the ESO telescopes two days before the *BeppoSAX* observation. We conclude that SAX J0103.2–7209 and 2E 0101.5–7225 are the same source, a relatively young and persistent X–ray pulsar in the SMC.

Subject headings: binaries: general — pulsars: individual (SAX J0103.2–7209; 2E 0101.5–7225) — stars: emission–line, Be — stars: rotation — X–ray: stars

1. INTRODUCTION

During 1997–1998 the number of X–ray pulsars found in the Small Magellanic Cloud (SMC) rapidly increased from three (SMC X–1, RX J0053.8–7226, and 2E 0050.1–7247; Lucke et al 1976; Hughes 1994; Israel et al. 1997) up 14 (for a review see Yokogawa et al. 1998) thanks to sensitive observations of the large area detectors on board the *RXTE* and *ASCA* satellites. The majority were found to be associated with massive Be spectral type stars showing intense H α emission lines. Only SMC X–1, which is associated with a super–giant B0 spectral type star in a 3.9 d orbital period binary system, is a persistent (although moderately variable) X–ray pulsar. For all the remaining X–ray pulsars pronounced variability (a factor ≥ 50) or, more often, transient behaviour has been definitively proven.

The source 2E 0101.5–7225 was detected at a nearly constant flux level in all the *Einstein*, *ROSAT* and *ASCA* pointings which surveyed the relevant region of the SMC (see Hughes & Smith 1994), but pulsations were not found due to poor statistics. Based on the accurate position obtained with the *ROSAT* HRI, these authors found that 2E 0101.5–7225 is very likely associated with a Be spectral–type star (R.A. = 01^h03^m13^s.86, Dec. = $-72^\circ 09' 14''.1$; equinox J2000). 2E 0101.5–7225 is located near the optical limb of the supernova remnant SNR 0101–72.4. Hughes & Smith (1994) present several arguments that make the Be/X–ray binary – SNR association unlikely.

In this *Letter* we report the discovery of 345 s pulsations from the source SAX J0103.2–7209 during a *BeppoSAX* observation of the SMC. The comparison with the data of past X–ray missions allows us to conclude that SAX J0103.2–7209 and 2E 0101.5–7225 are the same object, a persistent source with moderate variability (within a factor of 5–10). We also report the results of the timing analysis of a recent public *Chandra* observation and discuss optical observations carried out at ESO.

2. OBSERVATIONS AND DATA ANALYSIS**2.1. *BeppoSAX* observation**

The SMC field including the position of the 2E 0101.5–7225 was observed by the Narrow Field Instruments (NFIs) on board the *BeppoSAX* satellite (Boella et al. 1997a) on 1998 July 26–27 (effective exposure time of 40320 s). We used data from the Medium Energy (MECS; Boella et al. 1997b) and Low Energy (LECS; Parmar et al. 1997) instruments. A bright X–ray source ($\sim 3.7 \times 10^{-2}$ ct s^{–1}, 1–10 keV) was detected on–axis in the MECS, at R.A. = 01^h03^m13^s, Dec. = $-72^\circ 09' 16''$ (J2000; 90% confidence uncertainty radius of 30''). The MECS event list and spectrum were extracted from a circular region of 4' radius (corresponding to an encircled energy of $\sim 90\%$) around the X–ray position.

[†] The results reported in this Letter are partially based on observations carried out at ESO, La Silla, Chile (61.D–0513).

* Affiliated to the International Center for Relativistic Astrophysics (I.C.R.A.)




FIG. 1.— 1998 July (upper left panel) and 1999 August (lower left panel) power spectra of the 1–10 keV *BeppoSAX* MECS and *Chandra* ACIS light curves of 2E 0101.5–7225. The corresponding folded light curves at the best periods are also shown in the central panels. The *BeppoSAX* folded light curve is given separately in the 1–4 keV (S) and 4–10 keV (H) ranges. The period history of 2E 0101.5–7225 is shown in the right panel.




FIG. 2.— *BeppoSAX* (crosses) and *ROSAT* PSPC (triangles) spectra of 2E 0101.5–7225. The best fit model is shown, together with the corresponding residuals.

A 4' extraction radius ($\sim 85\%$ encircled energy) was also used for the LECS in order to minimize the contamination from the soft X-ray emission of the bright nearby source 2E 0102.3–7217. The local background was measured in a region of the MECS and LECS images far from any detected field sources.

The arrival times of the ~ 1500 photons were corrected to the barycenter of the solar system and background subtracted light curves were accumulated in 0.5 s bins. A single power spectrum was calculated over the entire time span covered by the observation in order to maximize the sensitivity to coherent pulsations. Significant power spectrum peaks were searched for using the algorithm described in Israel & Stella (1996). A highly significant peak ($\sim 9.4\sigma$ based on the fundamental only; see upper left panel of Fig. 1) was found at a frequency of 0.002889 Hz, corresponding to a period of 345.2 s. An accurate determination of the period was obtained by fitting the phases of the modulation over 4 different intervals of ~ 16000 s each. The scatter of the phase residuals was consistent with a strictly periodic modulation at the best period of 345.2 ± 0.3 s (90% confidence). A comparison of the 1–4 keV (panel S in Fig. 1) and 4–10 keV

(panel H) folded light curves provides marginal evidence for an energy dependent pulse profile: nearly sinusoidal at low energies, while double-horned above 4 keV. However the pulsed fraction (semi-amplitude of modulation divided by the mean source count rate) of $\sim 45\%$ is constant over the two energy intervals considered.

A spectral analysis was performed in the 0.7–6.5 and 1.6–10 keV energy ranges for the LECS and MECS, respectively. The spectra were rebinned to have at least 20 counts in each energy channel, such that χ^2 fitting techniques could be used. No single component model was found to fit the data well (a power-law gave a $\chi^2/dof = 41/15$; where dof is the degree of freedom). Among double component models, an absorbed power-law plus a black body gave the best fit ($\chi^2/dof = 15/15$) for a photon index of 1.0 ± 0.2 , $N_H \leq 3.8 \pm 3.8 \times 10^{21} \text{ cm}^{-2}$, and a black body temperature of 0.11 ± 0.03 keV (uncertainties refer to 1σ). The observed flux in the 2–10 keV energy band was $2.7 \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ (the soft component accounts for $\sim 15\%$ of the total) corresponding to an unabsorbed 2–10 keV X-ray luminosity of $1.2 \times 10^{36} \text{ erg s}^{-1}$ assuming a distance of 62 kpc (Laney & Stobie 1994).

2.2. Archival X-ray Observations

After the discovery of 345 s pulsations in 2E 0101.5–7225 (Israel et al. 1998), Yokogawa & Koyama (1998) found a signal at a period of 348.9 s during a 1996 May ASCA observation (absorbed luminosity of $5 \times 10^{35} \text{ erg s}^{-1}$; 2–10 keV) implying a period derivative (with respect to *BeppoSAX*) of -1.7 s yr^{-1} . The source was also detected by ASCA on 1993 May 12 and 1997 November 14 at an absorbed luminosity level of 6 and $5 \times 10^{35} \text{ erg s}^{-1}$, respectively. A re-analysis of the latter ASCA datasets allowed us to infer an upper limit (3σ confidence) in the 60–80% range on the pulsed fraction for a period of ~ 345 s.

To better constrain the absorption we used archival data from the *ROSAT* satellite (PSPC; 0.1–2 keV band) which observed the field of 2E 0101.5–7225 between 1991 October 8 and 1992 April 28 (sequence 600195; effective exposure time 26630 s). Assuming that the source showed the same flux and spectrum during the two observations, we fitted the *ROSAT* and *BeppoSAX* data together. We found that the two




FIG. 3.— $45'' \times 45''$ *Chandra* ACIS image of the region around 2E 0101.5–7225 (left panel). The image was smoothed with a Gaussian of $\sigma=2''$. The cross marks the optical position of the Be star identified by Hughes & Smith (1994). The double-peaked shape of the source is artificial (see text for details). $2' \times 2'$ $H\alpha$ image together with the *ROSAT* and *BeppoSAX* position circles (right panel). The proposed optical counterpart is also marked.

component model derived above again gave the best fit; a $\chi^2/dof = 23/25$ was obtained for a photon index of 1.1 ± 0.1 with N_H of $3.9 \pm_{2.4}^{4.0} \times 10^{21} \text{ cm}^{-2}$ and a black body temperature of $0.12 \pm_{0.03}^{0.04} \text{ keV}$ corresponding to an equivalent black body radius R_{bb} in the 2–60 km range (see Fig. 2; 1σ uncertainties). The corresponding power spectrum shows no significant (3σ confidence) peak in the 339–359 s period interval with a $\sim 60\%$ upper limit on the pulsed fraction.

2.3. *Chandra* observation

The field including 2E 0101.5–7225 was observed by the NASA Advanced X-ray Astrophysics Facility satellite *Chandra* on 1999 August 23 with the Imaging Spectrometer (ACIS; Garmire et al. 1992, Bautz et al. 1998 and references therein) in the high-resolution imaging mode for an effective exposure time of 19551 s. The source was detected at an off-axis angle of about $9'$ in the S4 (front illuminated) CCD of the ACIS-S detector with a count rate of $\sim 9 \times 10^{-2} \text{ ct s}^{-1}$ (in the 0.5–10 keV energy range; see Fig. 3 left panel). The source shows two emission peaks separated by $\sim 5''$. However the size of the emission region is comparable with the 90% encircled energy region of a source at $9'$ off-axis angle.

We extracted a 3.24 s binned light curve of the source from a region within $8''$ from the center of the emission: R.A. = $01^{\text{h}}03^{\text{m}}14^{\text{s}}.06$, Dec. = $-72^\circ 09' 15''.25$ (equinox J2000). The statistical uncertainty radius is only $\sim 0.5''$ but at this stage the uncertainty in the absolute positioning at such an off-axis angle might be as large as $90''$. However the detection of the pulsations clearly associates the *Chandra* source with that of *BeppoSAX*. We also note that the *Chandra* coordinates are consistent with the *ROSAT* HRI uncertainty circle and differ by less than $2''$ from those of the proposed optical counterpart (see below) making the association very likely. According to the *Chandra* source naming convention we designated it as CXO J010314.1–720915.

A single power spectrum was calculated over the entire observation. The search was performed over a period interval around that detected by *BeppoSAX* and assuming a maximum $|\dot{P}|$ of $\sim 3 \text{ s yr}^{-1}$ which translates into a search over only two Fourier frequencies (see Fig. 1 lower panels). A peak was de-

tected at a significance level of 7.5σ . A refined period was determined by means of the phase fitting technique; this gave a value of $343.5 \pm 0.5 \text{ s}$ (see Table 1). We note that the *Chandra* data were not corrected to the barycenter of the solar system. However, for a relatively long period pulsar in the direction of the SMC, the effect of the spacecraft and of the Earth motion, would cause a maximum correction of a factor of 10 smaller than the statistical uncertainty of the period given above. The pulse profile is sinusoidal (over the whole *Chandra* energy band) and the pulsed fraction is $\sim 45\%$ (see Fig. 1). This result implies that the pulsar is continuing to spin-up at a constant rate of -1.7 s yr^{-1} since 1996 (see Fig. 1; right panel).

In order to further address the issue of the double peaked *Chandra* image (see Fig. 3), we extracted two separate light curves for each of the two peaks. We found that the signal at 343.5 s was present with a similar pulsed fraction in both light curves; this result is consistent with the hypothesis of a point-like source. An independent confirmation was also obtained through a raytracing simulation, encompassing a monochromatic source (1.49 keV) and models for the mirror assembly and ACIS detector. The results indicate that expected PSF at an off-axis angle of $9'$ is artificially elongated in the direction connecting the two peaks detected in the *Chandra* image around the position of 2E 0101.5–7225; the two peaks are probably artefacts resulting from the small number of X-rays in the image. We conclude that the source is consistent with being point-like.

2.4. Optical Observations

Optical images ($H\alpha$ and $H\alpha$ red-continuum; 200 s each) of the *BeppoSAX* error circle of SAX J0103.2–7209 were obtained on 1998 July 24 at the Danish 1.5 m telescope with the Danish Faint Object Spectrometer Camera (DFOSC) at La Silla (Chile) in order to search for emission-line stars. The data were reduced using standard ESO-MIDAS procedures for bias subtraction, flat-field correction, aperture photometry and one dimensional stellar and sky spectra extraction. Within the *BeppoSAX* position circle we found only one $H\alpha$ active object: this is the O9–B1 III–Ve $m_V=14.8$ star originally suggested as the optical counterpart of 2E 0101.5–7225 (Hughes & Smith 1994). A 10 Å resolution 3800–8500 Å spectrum ($2''0$

TABLE 1
PERIOD HISTORY FOR 2E 0101.5–7225.

Mission	Instrument	Period (s)	Date	Exposure (s)	2–10 keV L_X ($10^{35} \text{ erg s}^{-1}$)	Reference
ASCA	GIS	348.9 ± 0.3	1996 May 21–23	32912	5	1
BeppoSAX	MECS	345.2 ± 0.2	1998 Jul 26–27	40320	11	2; this work
Chandra	ACIS	343.5 ± 0.5	1999 Aug 23	19551	—	this work

Note: Luminosities refer to absorbed fluxes. 90% uncertainties are reported.

1 – Yokogawa & Koyama (1998); 2 – Israel et al. (1998).

slit) of this star was obtained with the same instrument on 1998 October 20. Strong $H\alpha$ and $H\beta$ emission-lines were detected, with equivalent width $-20 \pm 2 \text{ \AA}$, and $-1.8 \pm 0.2 \text{ \AA}$, respectively. These results are in good agreement with those obtained by Hughes & Smith (1994) during observations carried out on December 1992.

3. DISCUSSION

Although 2E 0101.5–7225 shares several characteristics with other X–ray pulsators in the Magellanic Clouds (MCs) and in the Galactic plane (i.e., the spin period and pulsed fraction, the spectral shape at high energy and the presence of a soft thermal component, the association with a high mass companion, etc.), it has also some peculiar differences which make it unusual.

The absence of long-term variability greater than a factor of ~ 5 –10 over an interval of ~ 20 years is strongly suggestive of a *persistent* X–ray pulsar, the second in the SMC since the discovery of pulsations from SMC X–1. Moreover, the association of the companion with a Be spectral-type star makes 2E 0101.5–7225 the first example of a persistent (main sequence) Be/X–ray binary system in the SMC. 2E 0101.5–7225 is also a relatively low-luminosity (10^{35} – $10^{36} \text{ erg s}^{-1}$) X–ray pulsar. Finally, the inferred period derivative of -1.7 s yr^{-1} corresponds to a secular spin-up time-scale of $\sim 200 \text{ yr}$ which is the shortest of any known X–ray pulsar in a High Mass X–ray binary.

By using the P and $\dot{P}$ measurements and an X–ray luminosity in the 0.5 – $2 \times 10^{36} \text{ erg s}^{-1}$ range, we infer a magnetic field of 4 – $12 \times 10^{12} \text{ Gauss}$ and, correspondingly, a maximum magnetospheric radius of $r_m \sim 4 \times 10^9 \text{ cm}$ (see Lamb et al. 1973). Since the derived corotation radius r_{co} is $\sim 8 \times 10^9 \text{ cm}$, the r_m is considerably smaller than r_{co} , and accretion on the surface of the neutron star proceeds unaffected by the magnetosphere’s centrifugal drag, as long as the luminosity of the source is larger than $\sim 2 \times 10^{35} \text{ erg s}^{-1}$. Such a luminosity is not unusual for a bright X–ray pulsar in a nearly circular orbit around a giant OB companion star where the intense stellar wind con-

tinuously supplies matter to the compact object. Assuming a main sequence star, an orbital period of $\sim 300 \text{ d}$ is expected for 2E 0101.5–7225 based on the pulse period – orbital period correlation of Be star/X–ray pulsar binaries (Corbet 1984). A timing analysis of I band photometric measurements from the Optical Gravitational Lensing Experiment (OGLE) revealed no evidence of any coherent signal in the 1–50 d period interval (see Coe & Orosz 1999) suggesting that 2E 0101.5–7225 is a long period system.

The characteristics of 2E 0101.5–7225 have their closest analogy to those of the persistent low-luminosity (10^{35} – $10^{36} \text{ erg s}^{-1}$; 2–10 keV band) X–ray pulsar RX J0146.9+6121, a 1455 s (spin-up timescale of $\sim 250 \text{ yr}$) spinning neutron star in a binary system with a B1Ve companion star (see Reig et al. 1997; Haberl et al. 1998; Mereghetti et al. 2000). Two other recently discovered long period Be/X–ray pulsars in the SMC, namely AX J0051–733 ($P_s=323 \text{ s}$; Imanishi et al. 1999) and AX J0058–7203 ($P_s=280 \text{ s}$; Tsujimoto 1999), although highly variable (a factor of 10–100), have a maximum luminosity level close to that of 2E 0101.5–7225.

In conclusion, we discovered a new X–ray pulsar with a period of $\sim 345 \text{ s}$ in the SMC which is persistent, has a relatively low-luminosity, is rapidly spinning-up and is associated with a O9–B1 III–Ve star. The relatively high value inferred for the magnetic field and period derivative point to a young object. All these findings make 2E 0101.5–7225 an unusual X–ray pulsar which deserves more detailed investigations with instrumentation ranging from the X–ray to the IR band.

This work was partially supported through ASI and CNAAs grants. We thank the *Chandra* Data Archive (CDA) of the *Chandra* X–Ray Observatory Science Center (CXC) at CfA for a prompt release of the data. We thank R. Ragazzoni for obtaining the optical spectrum of the counterpart of 2E 0101.5–7225. We thank H. Tananbaum, R.J. Edgar and L. Burderi for their helpful comments. This work has been supported in part by NASA contract NAS8-39073.

REFERENCES

- Bautz, M. W., et al. 1998, SPIE, 3444, 210
 Boella, G., et al. 1997a, A&AS, 122, 299
 Boella, G., et al. 1997b, A&AS, 122, 327
 Coe, M.J. & Orosz, J.A. 1999, MNRAS, in press (astro-ph/9908303)
 Corbet, R.H.D. 1984, A&A, 141, 91
 Garmire, G. P. et al. 1992, AIAA, Space Programs and Technologies Conference, Huntsville, p. 8
 Haberl, F., Angelini, L. & Motch, C. 1998, A&A, 335, 587
 Hughes, J.P. 1994, ApJ, 427, L25
 Hughes, J.P. & Smith, R.C. 1994, PASP, 107, 1363
 Imanishi, K., et al. 1999, PASJ, in press (astro-ph/9910463)
 Israel, G.L. & Stella, L. 1996, ApJ, 468, 369
 Israel, G.L., et al. 1997, ApJ, 484, L141
 Israel, G.L., et al. 1998, IAU Circ. 6999
 Lamb, F.K., Pethick, C.J. & Pines, D. 1973, 184, 271
 Laney, C.D. & Stobie, R.S. 1994, MNRAS, 266, 441
 Lucke, R., et al. 1976, ApJ, 206, L25
 Mereghetti, S., Tiengo, A. & Israel, G.L. 2000, A&A, in press
 Parmar, A.N., et al. 1997, A&AS, 122, 309
 Reig, P., Fabregat, J. & Coe, M.J. 1997, A&A, 322, 193
 Tsujimoto, M., et al. 1999, PASJ, in press (astro-ph/9910505)
 Yokogawa, J. & Koyama, K. 1998, IAU Circ. 7009
 Yokogawa, J., et al. 1998, MPE Report 270, eds. Becker, W. & Itoh, M., p. 151